

Product no **AS06 172****Anti-PsaA | PSI-A core protein of photosystem I****Product information**

Immunogen	N-terminal part of recombinant PsaA protein from <i>Chlamydomonas reinhardtii</i> P12154
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	50 µl
Reconstitution	For reconstitution add 50 µl of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.
Additional information	PsaA is a hydrophobic protein and we recommend to use PVDF membrane for transfer to assure best results. This product can be sold containing ProClin if requested.

Application information

Recommended dilution	1 : 20 (IG), 1 : 1000-1 : 5000 (WB)
Expected apparent MW	82 55-60 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i> , <i>Begonia</i> sp., <i>Bryopsis corticulans</i> , <i>Chlamydomonas reinhardtii</i> , psychrophilic <i>Chlamydomonas</i> sp. UWO241 and <i>Chlamydomonas</i> sp. ICE-MDV, <i>Chlorella sorokiniana</i> , <i>Chlorella vulgaris</i> , <i>Chromochloris zofingiensis</i> , <i>Colobanthus quitensis</i> Kunt Bartl, <i>Craterostigma pumilum</i> , <i>Cytisus cantabricus</i> (Wilk.) Rchb. F., <i>Dianthus caryophyllus</i> , <i>Dioxonella giordanoi</i> (red alga), <i>Drosera capensis</i> , <i>Euonymus japonicus</i> , <i>Fraxinus ornus</i> , <i>Fucus vesiculosus</i> , <i>Haematococcus pluvialis</i> , <i>Halomicronema hongdechloris</i> , <i>Hieracium pilosella</i> L., <i>Hordeum vulgare</i> , <i>Lasallia hispanica</i> , <i>Nannochloropsis oceanica</i> strain IMET1, <i>Nicotiana benthamiana</i> , <i>Nicotiana tabacum</i> , <i>Oryza sativa</i> , <i>Pisum sativum</i> , <i>Marchantia polymorpha</i> (liverwort), micro <i>Nannochloropsis gaditana</i> , <i>Phaseolus vulgaris</i> , <i>Physcomitrium patens</i> , <i>Picea abies</i> , <i>Pinus strobus</i> , <i>Sinapsis alba</i> , <i>Spinacia oleracea</i> , <i>Synechococcus</i> PCC 7942, <i>Synechocystis</i> PCC 6803, <i>Syntrichia muralis</i> (Hedw.) Raab, <i>Scenedesmus obliquus</i> , <i>Tillandsia flabellate</i> , <i>Ulva prolifera</i>
Predicted reactivity	Algae, <i>Bigeloviella natans</i> , <i>Cannabis sativa</i> , <i>Catalpa bungei</i> , <i>Citrus x limon</i> , Cyanobacteria, <i>Cyanidioschyzon merolae</i> strain 10D, <i>Galdieria sulphuraria</i> , <i>Lycopersicon esculentum</i> , <i>Panax ginseng</i> , <i>Picea spinulosa</i> , <i>Pinus thunbergii</i> , <i>Phaeodactylum tricornutum</i> , <i>Populus alba</i> , <i>Thermosynechococcus elongatus</i> (strain BP-1), <i>Triticum aestivum</i> Species of your interest not listed? Contact us
Not reactive in	<i>Chromera velia</i>
Additional information	Immunogold localization has been done in leaf material of <i>Arabidopsis thaliana</i> .
Selected references	Penzler et al. (2024) . A pgr5 suppressor screen uncovers two distinct suppression mechanisms and links cytochrome b6f complex stability to PGR5. <i>Plant Cell</i> . 2024 Mar 27:koae098. doi: 10.1093/plcell/koae098. Mu et al. (2024) . Plastid HSP90C C-terminal extension region plays a regulatory role in chaperone activity and client binding. <i>Plant J</i> . 2024 Jul 5. doi: 10.1111/tbj.16917. Zhao et al. (2024) . Psb28 protein is indispensable for stable accumulation of PSII core complexes in Arabidopsis. <i>Plant J</i> . 2024 May 26. doi: 10.1111/tbj.16844. Kim et al. (2024) . Photoautotrophic cultivation of a <i>Chlamydomonas reinhardtii</i> mutant with zeaxanthin as the sole xanthophyll. <i>Biotechnol Biofuels Bioprod</i> . 2024 Mar 14;17(1):41. doi: 10.1186/s13068-024-02483-8. Khaig and Eaton-Rye (2023) . Lys264 of the D2 Protein Performs a Dual Role in Photosystem II Modifying Assembly and Electron Transfer through the Quinone-Iron Acceptor Complex. <i>Biochemistry</i> 2023, 62, 18, 2738–2750 Okegawa et al. (2023) . x- and y-type thioredoxins maintain redox homeostasis on photosystem I acceptor side under fluctuating light. <i>Plant Physiol</i> . 2023 Nov 22;193(4):2498-2512. doi: 10.1093/plphys/kiad466. Sulli et al. (2023) . Generation and physiological characterization of genome edited <i>Nicotiana benthamiana</i> plants containing zeaxanthin as the only leaf xanthophyll. <i>Planta</i> . 2023 Oct 5;258(5):93. doi: 10.1007/s00425-023-04248-3. Jiang et al. (2023) . Toxic effects of lanthanum (III) on photosynthetic performance of rice seedlings: Combined chlorophyll fluorescence, chloroplast structure and thylakoid membrane protein assessment. <i>Ecotoxicol Environ Saf</i> . 2023 Nov 15:267:115627. doi: 10.1016/j.ecoenv.2023.115627. Kafri et al. (2023) . Systematic identification and characterization of genes in the regulation and biogenesis of photosynthetic machinery. <i>Cell</i> . 2023 Dec 7;186(25):5638-5655.e25. doi: 10.1016/j.cell.2023.11.007.


[Nagy et al. \(2023\)](#). Photoautotrophic and sustained H₂ production by the pgr5 mutant of *Chlamydomonas reinhardtii* in simulated daily light conditions. *International Journal of Hydrogen Energy* Volume 53, 31 January 2024, Pages 760-769.

[Hu et al. \(2023\)](#). Drought affects both photosystems in *Arabidopsis thaliana*. *New Phytol.* 2023 Oct;240(2):663-675. doi: 10.1111/nph.19171. Epub 2023 Aug 2.

[Ivanov et al. \(2022\)](#) The decreased PG content of pgp1 inhibits PSI photochemistry and limits reaction center and light-harvesting polypeptide accumulation in response to cold acclimation. *Planta* 255, 36 (2022). <https://doi.org/10.1007/s00425-022-03819-0>

[Lim et al \(2022\)](#). *Arabidopsis* guard cell chloroplasts import cytosolic ATP for starch turnover and stomatal opening. *Nat Commun.* 2022 Feb 3;13(1):652. doi: 10.1038/s41467-022-28263-2. PMID: 35115512; PMCID: PMC8814037.

[Lim et al \(2022\)](#) *Arabidopsis* guard cell chloroplasts import cytosolic ATP for starch turnover and stomatal opening. *Nat Commun.* 2022 Feb 3;13(1):652. doi: 10.1038/s41467-022-28263-2. PMID: 35115512; PMCID: PMC8814037.



2 µg of total protein from **(1)** *Arabidopsis thaliana* leaf, **(2)** *Hordeum vulgare* leaf, **(3)** *Chlamydomonas reinhardtii* total cell, **(4)** *Synechococcus* sp. 7942 total cell all extracted with Protein Extraction Buffer, PEB ([AS08 300](#)), were separated on **4-12%** NuPage (Invitrogen) **LDS-PAGE** and blotted 1h to **PVDF**. Blots were blocked immediately following transfer in 2% blocking reagent in 20 mM Tris, 137 mM sodium chloride pH 7.6 with 0.1% (v/v) Tween-20 (TBS-T) for 1h at room temperature with agitation. Blots were incubated in the primary antibody at a dilution of 1: 10 000 for 1h at room temperature with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed once for 15 min and 3 times for 5 min in TBS-T at room temperature with agitation. Blots were incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, recommended secondary antibody [AS09 602](#)) diluted to 1:50 000 in 2% blocking solution for 1h at room temperature with agitation. The blots were washed as above and developed for 5 min with chemiluminescence detection reagent according the manufacturers instructions. Images of the blots were obtained using a CCD imager (FluorSMax, Bio-Rad) and Quantity One software (Bio-Rad). Exposure time was 10 seconds.